

United States Patent [19]

Akhavan-Tafti

[11] Patent Number:

6,001,614

[45] Date of Patent:

*Dec. 14, 1999

[54] METHODS OF SYNTHESIZING LABELED POLYNUCLEOTIDES BY LIGATION OF MULTIPLE OLIGOMERS

[75] Inventor: Hashem Akhavan-Tafti, Howell, Mich.

[73] Assignee: Lumigen, Inc., Southfield, Mich.

[*] Notice: This patent is subject to a terminal dis-

claimer.

[21] Appl. No.: 09/241,353

[22] Filed: Feb. 2, 1999

Related U.S. Application Data

[63]	Continuation	of application	No.	09/121,887,	Jul.	24,	1998.

[51] Int. Cl.⁶ C12P 19/34; C12Q 1/68; C07H 21/04

[56] References Cited

U.S. PATENT DOCUMENTS

5,478,724 5,663,062 5,695,933 5,750,341 5,770,367	12/1995 9/1997 12/1997 5/1998 6/1998	Brennan 435/6 Morse 435/5 Sorge 435/91.1 Schalling 435/91.52 Macevicz 435/6 Southern 435/6 Yager 435/6
5,888,731	3/1999	Yager 435/6

FOREIGN PATENT DOCUMENTS

88311741 6/1989 European Pat. Off. .

OTHER PUBLICATIONS

S. Dubiley, E. Kirilov, Y. Lysov, A. Mirzabekov Nucl. Acids Res., 25, 2259–2265 (1997).
R. Handley, H. Akhavan–Tafti, A. P. Schaap, J. Clin. Ligand Assay, 20(4) 302–312 (1997).

K.D.James, A.D.Ellington, Chemistry & Biology 4, 595-605 (1997).

T.Kaczorowski, W.Szybalski, Gene 179, 189-193 (1996).

L.E.Kotler, D. Zevin-Sonkin, I.A. Sobolev, A.D. Beskin L.E. Ulanovsky, Proc. Natl. Acad. Sci. USA, 90, 4241–4245 (1993).

T.Li, K.C. Nicolaou, Nature, 369, 218-221 (1994).

C.E.Pritchard, E.M.Southern, Nucl. Acids Res., 25, 3403–3407 (1997).

D.Sievers, G. Von Kiedrowski, Nature, 369, 221-224 (1994).

W.Stemmer, A.Crameri, K.D.Ha, T.M.Brennan, H.L. Heyneker, Gene, 164, 49-53 (1995).

D.Y.Wu, R.B. Wallace, Genomics, 4,560-569 (1989).

Primary Examiner—Carla J. Myers
Assistant Examiner—Diana Johannsen
Attorney, Agent, or Firm—Richard S. Handley

57] ABSTRACT

Methods of synthesizing polynucleotides are disclosed involving the simultaneous ligation of a set of oligomer 5'-phosphates onto a template-bound primer. The set of these oligomers can be preselected to contain oligomers which are complementary to the template strand or the oligomers can be supplied as a library and allowed to self select. The synthesis by ligation can proceed unidirectionally or bidirectionally from the primer and can be used to synthesize both strands simultaneously by the use of two primers. The ligation is preferably performed with a ligase enzyme. The methods of synthesis are useful in a variety of applications, including cloning, amplification, labeling, diagnostic assays, mutation analysis and screening, gene expression monitoring and sequence analysis.

23 Claims, 6 Drawing Sheets

